

TOWARDS
PHYSICAL REALIZATION
OF
TOPOLOGICALLY PROTECTED
QUBITS

With the collaboration of:

Lev Ioffe, Misha Gershenson and his experimental group
(Rutgers)

Wiebke Guichard, Olivier Buisson (Grenoble)


Thomas Coudreau, Perola Milman (MPQ)

Uri Gavish (LPTHE)


Lattice gauge theory in deconfined regime I

A. Kitaev, Ann. Phys. **303**, 2, (2003)

$$Z_2 \text{ charge } U_i = \prod_j^{(i)} \sigma_{ij}^x$$



$$Z_2 \text{ flux } B_{\square} = \prod_{ij \in \square} \sigma_{ij}^z$$



$$H_{\text{Kitaev}} = -\frac{\Delta_c}{2} \sum_i U_i - \frac{\Delta_f}{2} \sum_{\square} B_{\square}$$

Localized excitations with finite energy gap

Ground-state degeneracy depends on global topology of the lattice.

Lattice gauge theory in deconfined regime II

Two-fold degenerate ground-state on a cylinder

Degeneracy enforced by non-local symmetries:



row operators:

$$P_i = \prod_j \sigma_{ij}^z$$

column operators:


$$Q_j = \prod_i \sigma_{ij}^x$$

$$\{P_i, Q_j\} = 0$$




The only dangerous errors are non-local !
 They are suppressed by a factor $(\text{noise}/\Delta_{c,f})^L$

Electric noise transfers one Z_2 flux along v-path and flips P_i : Relaxation in *flux* basis or dephasing in *charge* basis.



Magnetic noise transfers one Z_2 charge along h-path and flips Q_j : Relaxation in *charge* basis or dephasing in *flux* basis.



Basics of Josephson junction arrays

ϕ_j ; local phase of Cooper pair condensate


$\hat{n}_j = \frac{\partial}{i\partial\phi_j}$: number of Cooper pairs on island j

$$\Delta\phi_j \Delta n_j \simeq 2\pi$$

$$A_{ij} = \frac{2\pi}{\Phi_0} \int_i^j \vec{A}_{ij} \cdot d\vec{r}$$

$$H = -E_J \sum_{\langle ij \rangle} \cos(\phi_i - \phi_j - A_{ij}) + \frac{E_C}{2} \sum_{ij} (C_{ij}^{-1}) \hat{n}_i \hat{n}_j$$

E_J : Josephson coupling energy



E_C : Charging energy


A rhombus with half a flux quantum

Define $\theta_{ij} = \phi_i - \phi_j - A_{ij}$, then:

$$\theta_{12} + \theta_{23} + \theta_{34} + \theta_{41} \equiv \pi, \text{ mod } 2\pi$$

→ Get two-fold degenerate classical ground-state, with $\theta_{ij} = \pm\frac{\pi}{4}$

→ Quantum fluctuations ($E_C \neq 0$) of phases lift this degeneracy




Principle of a protected qubit

$$H = -\frac{\Delta_f}{2} \sum_{i=1}^M \tau_i^z \tau_{i+1}^z - \frac{\Delta_c}{2} \sum_{i,j,j'} \sigma_{i,j}^x \sigma_{i,j'}^x$$

Very similar to the Kitaev model:

$$\tau_i^z = \prod_{j=1}^N \sigma_{i,j}^z = P_i$$

Josephson array version



$\sigma_{i,j}^z$: chirality of rhombus j
 $(1 \leq j \leq N)$ in chain i
 $(1 \leq i \leq M)$

Degenerate ground-states of a single chain

$$H_i = -\frac{\Delta c}{2} \sum_{j,j'} \sigma_{i,j}^x \sigma_{i,j'}^x$$

Define $|\pm\rangle_{i,j} = (|\uparrow\rangle_{i,j} \pm |\downarrow\rangle_{i,j}) / \sqrt{2}$.

The two ground-states of H_i are $|+, +, \dots, +\rangle_i$ and $|-, -, \dots, -\rangle_i$

$$[H_i, \prod_{j=1}^N \sigma_{i,j}^z] = 0$$

Recall the definition: $\tau_i^z = \prod_{j=1}^N \sigma_{i,j}^z$

$$\begin{aligned} |\tau_i^z = \uparrow\rangle_i &= (|+, +, \dots, +\rangle_i + |-, -, \dots, -\rangle_i) / \sqrt{2} \\ |\tau_i^z = \downarrow\rangle_i &= (|+, +, \dots, +\rangle_i - |-, -, \dots, -\rangle_i) / \sqrt{2} \end{aligned}$$

Ground-states of a single chain($N = 4$)

$$|\tau_i^z = \uparrow\rangle_i \propto |+, +, +, +\rangle_i + |-, -, -, -\rangle_i \propto \begin{aligned} &|\uparrow\uparrow\uparrow\uparrow\rangle_i + |\uparrow\uparrow\downarrow\downarrow\rangle_i + |\uparrow\downarrow\uparrow\downarrow\rangle_i + \\ &|\uparrow\downarrow\downarrow\uparrow\rangle_i + |\downarrow\uparrow\uparrow\downarrow\rangle_i + |\downarrow\uparrow\downarrow\uparrow\rangle_i + \\ &|\downarrow\downarrow\uparrow\uparrow\rangle_i + |\downarrow\downarrow\downarrow\downarrow\rangle_i \end{aligned}$$

$$|\tau_i^z = \downarrow\rangle_i \propto |+, +, +, +\rangle_i - |-, -, -, -\rangle_i \propto \begin{aligned} &|\uparrow\uparrow\uparrow\downarrow\rangle_i + |\uparrow\uparrow\downarrow\uparrow\rangle_i + |\uparrow\downarrow\uparrow\uparrow\rangle_i + \\ &|\downarrow\uparrow\uparrow\uparrow\rangle_i + |\uparrow\downarrow\downarrow\downarrow\rangle_i + |\downarrow\uparrow\downarrow\downarrow\rangle_i + \\ &|\downarrow\downarrow\uparrow\downarrow\rangle_i + |\downarrow\downarrow\downarrow\uparrow\rangle_j \end{aligned}$$

τ_i^z is associated to the superconducting phase difference across line i of the array, which is twofold degenerate.

Protection of local τ_i^z against decoherence


Noise of the form $\sum_{i,j} h_{i,j}^x(t) \sigma_{i,j}^x$ flips τ_i^z , but the strongly coupled M parallel chains correct it efficiently: such error costs an energy of the order E_J/N .

Noise of the form: $\sum_{i,j} h_{i,j}^z(t) \sigma_{i,j}^z$ commutes with τ_i^z , or equivalently, induces a term of the form:

$$|+, +, \dots, +\rangle_i \langle -, -, \dots, -|_i + h.c.$$

But this decoherence can be made very small !

Assume static and uniform noise $h_{i,j}^z(t) = h$




Ground-state doublet splitting $\simeq h^z \left(\frac{h^z}{\Delta_c} \right)^{N-1}$

So, if $h^z / \Delta_c < 1$, spin flip induces only exponentially small dephasing when N is large !

Same conclusion with non-uniform and dynamical noise $h_{i,j}^z(t)$, provided $\omega_{\text{noise}} \ll \Delta_c$

Experimental design



Total phase γ across the array is fixed by the combined effect of the large junction and the flux through the big loop. Qubit encoded in the phase (0 or π) of the **central island**.

Experimental diagnosis of topological protection

$$E_{\text{array}} = -E_1(\Phi_{\text{Rh}}) \cos \gamma - E_2(\Phi_{\text{Rh}}) \cos 2\gamma$$

If $\Phi_{\text{Rh}} \rightarrow \Phi_0(n + 1/2)$, n integer, then $E_1 \rightarrow 0$:

$\Phi_0/2$ periodicity,

else, $E_1 > E_2$: Φ_0 periodicity





$$\Delta I_c = \frac{2e}{\hbar} E_1 \sin \gamma + \frac{4e}{\hbar} \sin 2\gamma$$

Manifestation of protection: for $\Phi_{\text{Rh}} = \Phi_0(n + 1/2) + \Delta\Phi_{\text{Rh}}$,


$$E_1 \simeq E_J \left(\frac{E_J}{\Delta_c} \right)^{N-1} (\Delta\Phi_{\text{Rh}})^N$$

Exponential improvement with system length !


Experimental realization: M. Gershenson et al. (2007)




Evidence for finite Δ_c and charge $4e$ condensate



Phase stiffness E_2 of charge $4e$ condensate




Measuring the Z_2 charge of the central island



Critical current of the array depends on the Z_2 charge of the central island:

Provides a way to measure τ^x


Manipulating the protected qubit



Remarks about conserved quantities in the real physical system

If local magnetic fluxes are exactly $\Phi_0/2$ for each rhombus: local Z_2 charge is conserved, which implies $[Q_j, H_{\text{array}}] = 0$.


The array model allows for quantum tunneling processes where Z_2 vortices are created in pairs, which lead to $[P_i, H_{\text{array}}] \neq 0$. If we can neglect the processes where a single Z_2 vortex crosses the whole system, we can map the array into an **effective** Kitaev model where $[P_i, H_{\text{effective}}] = 0$.



Effect of noise during the τ^z manipulation

$$\Delta H(t) = f(t) \sum_{i,j} \sigma_{ij}^z + \sum_a \sum_{i,j} \epsilon_{ij}^a(t) \sigma_{ij}^a$$

σ_{ij}^z flips Q_j
 σ_{ij}^x flips P_i
 σ_{ij}^y flips P_i and Q_j



$$P_i = \prod_j \sigma_{ij}^z$$

$$Q_j = \prod_i \sigma_{ij}^x$$

$$\{P_i, Q_j\} = 0$$

Effect of various perturbations in qubit space

	Q_j 's conserved	Q_j 's flip
P_i 's conserved	$\mathbf{1}$	τ^z
P_i 's flip	τ^x	τ^y

Noise coupled to σ_{ij}^z

	Q_j 's conserved	Q_j 's flip	
P_i 's conserved	$(f + \epsilon^z)^n \mathbf{1}$	$(f + \epsilon^z)^{N+n} \tau^z$	n even
P_i 's flip	$0 \tau^x$	$0 \tau^y$	

Relative effect is of order $\frac{\epsilon^z}{f}$: the angle of the qubit rotation is not very robust.

Noise coupled to σ_{ij}^x

	Q_j 's conserved	Q_j 's flip	
P_i 's conserved	$f^n (\epsilon^x)^m \mathbf{1}$	$f^{N+n} (\epsilon^x)^m \tau^z$	m, n even
P_i 's flip	$f^n (\epsilon^x)^{M+m} \tau^x$	$f^{N+n} (\epsilon^x)^{M+m} \tau^y$	

Relative effect on rotation angle is of order $(\frac{\epsilon^x}{\Delta})^2$.

Relative size of τ^x term is of order $(\frac{\epsilon^x}{\Delta})^M (\frac{\Delta}{f})^N$.

Relative size of τ^y term is of order $(\frac{\epsilon^x}{\Delta})^M$.

Noise coupled to σ_{ij}^y

	Q_j 's conserved	Q_j 's flip
P_i 's conserved	$f^p (\epsilon^y)^m \mathbf{1}$	$f^{N-m+p} (\epsilon^y)^m \tau^z$
P_i 's flip	$f^n (\epsilon^y)^{M+m} \tau^x$	$f^{N-M+p} (\epsilon^y)^{M+m} \tau^y$

$m, p, n - M$, even

Relative effect on rotation angle is of order $(\frac{\epsilon^x}{f})^2$.

Relative size of τ^x term is of order $(\frac{\epsilon^x}{f})^N$ (if $M = N$).

Relative size of τ^y term is of order $(\frac{\epsilon^x}{f})^N$ (if $M = N$).

Remarks about the τ^x manipulation

$$\Delta H(t) = g(t) \sum_{i,j} \sigma_{ij}^z + \sum_a \sum_{i,j} \epsilon_{ij}^a(t) \sigma_{ij}^a$$

Formally similar to the τ^z manipulation !


Summary: effect of noise during qubit manipulation

Angle of rotation is no longer robust: the strength of the manipulation term should be larger than the noise which acts in the same channel.

Provided this condition is satisfied, there is still **some form of protection** for the **direction** of the rotation applied in qubit space, in the sense that a **larger size** helps.

Open questions on the effect of noise during qubit manipulation

i) Putting back the true array Hamiltonian, for which strictly speaking, $[P_i, H_{\text{array}}] \neq 0$. As we have seen, these terms are associated to processes of the form:



which lift the degeneracy of the **central island**, whose **phase** tunnels between 0 and π , thus selecting its **Z_2 charge** as the good quantum number of the **qubit**.

ii) Include explicitly dynamical noise during manipulation


Computational issues I: hierarchical approximation

Series composition of Z_2 junctions

$$V(\phi) = -E_2 \cos(2\phi)$$

$$E'_2 = \left[1 - \frac{7}{256} \left(\frac{E_2}{E_C} \right)^2 \right] \frac{1}{8} \frac{E_2^2}{E_C}$$



$$E'_C = \left[1 - \frac{1}{16} \left(\frac{E_2}{E_C} \right)^2 \right] 2E_C$$



Parallel composition


$$E'_2 = K E_2$$

$$E'_C = K^{-1} E_C$$




Computational issues II: single rhombus as Z_2 junction



Effective E_2



Test of coarse graining:
2 and 3 rhombi in series



Effective capacitance



Conclusions

- 1) Kitaev's Z_2 lattice model implemented in the low energy sector of some Josephson junction arrays.
- 2) These arrays are composed of fully frustrated rhombi.
- 3) **Topological protection** arises in the phase where quantum phase fluctuations destroy the $2e$ condensate, while preserving the $4e$ condensate.
- 4) Experimental evidence for this phase: observation of enhanced immunity against **static** flux fluctuations, evidence of a finite Δ_c .
- 5) Some form of protection is still expected to hold during qubit manipulation.